

CellME Berlin 2023

International Forum on Cell Manufacturing & Engineering

7-8 November 2023. Berlin, Germany

www.cellme.de

Organized by:

DEMACH Event- und Veranstaltungs GmbH
Berlin, Germany



TARGET YOUR FOCUS VERTICALS

Check out our Upcoming Events
www.demach-events.de



ATTENDEE ACQUISITION

Hosting your own event means you have full control over the entire event experience, including the branding, promotions, content, and guest list.

You focus on content and programming, let us worry about the attendee list. Our team and technology deliver targeted, personalized event invitation message to your target prospects via most effective marketing channels. Our unique approach allows us to guarantee 100% of our leads to be fresh, up to date, and matching your ideal attendee profile, we will find the prospects and build a campaign that drives them to action.

DEMACH

Events for Innovation

EVENT MARKETING SERVICE



TAILORED
THE EVENT
TO YOUR GOAL



HOST YOUR EXCLUSIVE EVENT WITH US

PRODUCT DEMONSTRATION | SEMINAR | WORKSHOP/TUTORIAL | PRODUCT TRAINING
TECH DAY | USER CONFERENCE | BUSINESS/CHANNEL PARTNER EVENT

Want to learn more or request a pricing quote?

Please contact us at guestevent@demach-events.de and our service team will get in touch.


INDEX

EXHIBITOR

- ACCELERATE GMBH / GERMANY
- CELLBOX SOLUTIONS GMBH / GERMANY
- DON WHITLEY SCIENTIFIC LTD / UNITED KINGDOM
- FACELLITATE GMBH / GERMANY
- FIBERCELL SYSTEMS INC. / UNITED STATES
- HISS DIAGNOSTICS GMBH / GERMANY
- NEST SCIENTIFIC EUROPE B.V. / THE NETHERLANDS
- OMNI LIFE SCIENCE GMBH & CO. KG / GERMANY
- PHIO SCIENTIFIC GMBH / GERMANY
- SHANGHAI RUIYU BIOTECH CO., LTD. (PART OF ALIT BIOTECH / P.R.CHINA
- SPHERE FLUIDICS LIMITED / UNITED KINGDOM
- VITAL3D TECHNOLOGIES, UAB / LITHUANIA
- WORLD PRECISION INSTRUMENTS GERMANY GMBH / GERMANY
- XCELTIS GMBH / GERMANY

SPEAKER

- CELLBOX SOLUTIONS / GERMANY
- FACELLITATE GMBH / GERMANY
- FIBERCELL SYSTEMS INC / UNITED STATES
- FREIE UNIVERSITÄT BERLIN / GERMANY
- INSTITUTE OF MUSCULOSKELETAL MEDICINE, UNIVERSITY OF MUENSTER / GERMANY
- LUDWIG-MAXIMILIANS-UNIVERSITY MUNICH, PHIO SCIENTIFIC GMBH / GERMANY
- MEDICAL UNIVERSITY OF GRAZ / AUSTRIA
- ORGONEX / THE NETHERLANDS
- OXFORD OPTRONIX LTD. / UNITED KINGDOM
- SPHERE FLUIDICS LTD. / UNITED KINGDOM
- STONY BROOK CANCER CENTER, STONY BROOK UNIVERSITY, NEW YORK / UNITED STATES
- SUDHIN BIOPHARMA / UNITED STATES
- THE UNIVERSITY OF MANCHESTER / UNITED KINGDOM
- UNIVERSITÄTSKLINIKUM WÜRZBURG - FRAUNHOFER ISC / GERMANY
- VITAL3D TECHNOLOGIES / LITHUANIA
- WARSAW UNIVERSITY OF TECHNOLOGY / POLAND

 Last updated: November 03, 2023

LEGAL INFORMATION

Liability for Contents

The content of our pages has been compiled with utmost care. However, we do not assume any liability concerning the correctness, completeness and topicality of their content. In accordance with Article 7, Section 1 of the TMG (German Telemedia Act) we, as a service provider, are responsible for our own content on these pages in compliance with the general law. According to paragraphs 8 to 10 of the TMG, however, we are not obliged either to monitor third party information, which has been transferred or stored, or to search for circumstances that would indicate illegal activity. The responsibility to remove or block the use of information according to the general law shall remain unaffected. Any liability in this respect, however, is only possible from the moment at which knowledge of a specific violation of the law is obtained. We shall immediately remove this content upon discovery of such infringements.

Liability for Links

Our offer contains links to external websites of third parties, on the content of which we do not have any influence. Therefore, we cannot assume liability for this content. The liability for linked pages lies exclusively with their operators or providers. The linked pages were checked for possible legal violations at the time the links were created. Illegal content could not be identified at the time of setting the links. However, we cannot reasonably be expected to constantly monitor the content of the linked pages without any indication that the law has been violated. We shall immediately remove such links upon discovery of infringements.


Copyright

The content and works on these pages compiled by their operators are subject to German copyright law. Duplication, processing, distribution as well as any kind of exploitation, which lies beyond the limits of copyright law, requires the written consent of the respective author or originator. Downloads and copies from this page are only permitted for private, not commercial use. Insofar as the content of this page was not compiled by the operator, third party copyrights shall be observed. In particular, third party content shall be marked as such. In the event that you still become aware of a copyright violation, we kindly ask you to inform us accordingly. We shall immediately remove such content upon discovery of infringements.


COUNTRIES OF ATTENDEES

CellME Berlin 7.-8.11. 2023. Berlin, Germany

Last updated: 3 November 2023. 16:00 CET



CellME Berlin 2023 Conference Schedule

 The conference language is English





Day1: Tuesday, November 7, 2023	TIME	TOPIC	PRESENTER	COUNTRY
	10:00	<i>Opening Remarks</i>		
	10:10	Peptigels: a technological platform for the design of fully defined synthetic 3d hydrogel scaffolds for cell engineering applications.	THE UNIVERSITY OF MANCHESTER	UNITED KINGDOM
	10:35	Agrostemmoside e: a potent tool for transfection assays	FREIE UNIVERSITÄT BERLIN	GERMANY
	10:50	Spheroid culture of murine primary chondrocytes and atdc5 chondrocyte cell line for musculoskeletal research	INSTITUTE OF MUSCULOSKELETAL MEDICINE, UNIVERSITY OF MÜNSTER	GERMANY
	11:05	<i>Coffee Break</i>		
	11:35	Development of an organoid-based cell transformation assay for intestinal carcinogenicity screening of chemicals	STONY BROOK CANCER CENTER, STONY BROOK UNIVERSITY, NEW YORK	UNITED STATES
	11:55	3-D hollow fiber perfusion bioreactors for recapitulation of the in vivo microenvironment: a better way to grow cells.	FIBERCELL SYSTEMS INC	UNITED STATES
	12:25	<i>Lunch Time</i>		
	14:00	Cellbox – the shipment solution for living cells, organoids and sensitive probes.	CELLBOX SOLUTIONS	GERMANY
	14:30	Novel automated picodroplet-based technology to streamline cell line development and cell engineering workflows	SPHERE FLUIDICS LTD.	UNITED KINGDOM
	15:00	Vital light 3d – towards next breakthrough in bioprinting	VITAL3D TECHNOLOGIES	LITHUANIA
	15:25	New non-invasive, label-free monitoring approach for 2D and 3D cell culture	LUDWIG-MAXIMILIANS-UNIVERSITY MUNICH, PHIO SCIENTIFIC GMBH	GERMANY
	15:50	Networking		
	17:00	<i>End</i>		

Day2: Wednesday, November 8, 2023	TIME	TOPIC	PRESENTER	COUNTRY
	9:30	<i>Opening Remarks</i>		
	9:40	Creation of skin cell spheres and improved methods for measuring cell viability in the 3D model	WARSAW UNIVERSITY OF TECHNOLOGY	POLAND
	10:05	AI-powered analysis in robot-assisted three-dimensional tissue models production	UNIVERSITÄTSKLINIKUM WÜRZBURG - FRAUNHOFER ISC	GERMANY
	10:30	<i>Coffee Break</i>		
	11:00	Single use disposable inclined settling bioreactor for gentle expansion, concentration & harvest of cell and gene therapies	SUDHIN BIOPHARMA	UNITED STATES
	11:25	RPMotion - A novel bioreactor for rapid organoid production	ORGONEX	THE NETHERLANDS
	11:55	<i>Lunch Time</i>		
	13:30	Animal-free and chemically defined coating solution for biologically relevant in vitro assays	FACELLITATE GMBH	GERMANY
	14:00	From microenvironments to automation: supporting reproducibility of cell manufacturing and organoid research	OXFORD OPTRONIX LTD.	UNITED KINGDOM
	14:30	Patient-derived cancer cell models	MEDICAL UNIVERSITY OF GRAZ	AUSTRIA
	14:55	Networking		
	16:30	<i>End</i>		

/Be aware that the start and end times for each session are tentative. /This schedule is subject to change without prior notice. Check back for updates on www.cellme.de. (Lasted updated: 03.11.2023)



Organize your cryostocks



www.cellseeker.org

INVENTORY


Create a virtual inventory and track the location of every single vial in your cell collection.

CELLS

Create data sheets of your cell lines and link the information to your cell stocks.

Cellseeker GmbH

support@cellseeker.org
+49 160 624 060 7
Osterfeldstraße 12- 14
22529 Hamburg
Germany



Manage your lab's cell collection.

Assign users and permissions.


Give selected access to external partners.



Review the quality status of your cell stocks.

Control and track cell usage.

Trace back history of cell banks.



Track location and movement of your stocks.

Check-in & check-out vials from a virtual tank.

Import existing databases.

THE NEW GENERATION FOR LIVE CELL SHIPPERS



CELLBOX
Flight 2.0



Made in Germany

The **Cellbox Flight 2.0** makes it possible to perform long distance transport by air. It was uniquely engineered to use dry ice as a CO₂ source to comply with airline regulations. The user-programmable CO₂ and temperature values allow you to optimize the conditions in the incubation chamber for your precious biological material.

Benefits of live cell transport



- ➔ No pH-shift or use of metabolically active cryoprotectants
- ➔ No altered protein expression; no mechanic or osmotic stress
- ➔ Intelligently maintains internal environment for up to 32 hours
- ➔ High cell viability and accurate continuous data logging
- ➔ No freezing, no thawing - Cells arrive ready-to-use



CO₂ SUPPLY

Provides regulated CO₂ environment



TEMPERATURE

Maintains temperature between 28-38°C



APP

Data logging and export via Cellbox App



RECHARGEABLE

Li-ION battery with external 100 - 230V power supply

CONTACT

+49 (0) 40 226 316 4-10 enquiries@cellbox-solutions.com www.cellbox-solutions.com
Englische Planke 8 20459 Hamburg | An der Bahn 5 22844 Norderstedt | Germany



Whitley H35 Hypoxystation

- ✓ Precise control of O₂ and CO₂
 - ✓ Precise control of temperature and humidity
 - ✓ Colour touchscreen control panel for ease of use
 - ✓ Easy, accurate, 2-point oxygen sensor calibration
 - ✓ Integral 12 litre airlock
 - ✓ HEPA option available
-

Reliable cell-based assays are urgently needed to develop predictable model and testing systems

BIOFLOAT™

Current challenges

- Model systems mimicking cell natural microenvironment are required
- Rapid and reliable generation of disease models is needed for translational research
- An animal-free environment for human cell cultivation is preferred
- Optimal assay systems for stem cell or cancer research, pharmacology or toxicology are urgently needed

Our solution

- Rapid and round spheroid generation allowing cell-cell contacts to resemble natural environment
- Cell and protein attachment reliably avoided for optimal cell analysis
- Applicable to various complex systems, including bioreactors and microfluidic devices


BIOFLOAT™ outperform current competitors




BIOFLOAT™ – features

- DIY coating of polystyrene, glass and PDMS by simple rinsing process – no chemical treatments
- Stable anti-adhesive layer
- Coating prevents attachment of proteins and cells
- Generation of cell instructive surfaces by biological tags

BIOFLOAT™ – spheroids with challenging cells



- Round uniform hepatocytes spheroids. No satellites
- CYP activity measurable -> vital hepatocytes spheroids
- Human hepatocytes are highly relevant models for liver diseases/ damage in toxicological studies



- Round and reproducible spheroids. No satellites
- Rapid and reliable formation of vital spheroids within few hours. No centrifugation needed
- FACILITATES the development of *in vivo* similar model and testing systems for research and industry application



BIOFLOAT™
Order your free sample



faCellitate GmbH
Franz-Volhard Straße 5,
68167 Mannheim
Germany


www.facellitate.com

info@facellitate.com

Contact: valentina.fermi@facellitate.com

Hollow Fiber Bioreactors: A Better Way to Grow Cells

A closer look at the FiberCell Systems cartridge



APPLICATIONS



3-D Cell Culture

The medium-sized cartridge offers 3,000 cm² of surface area, equivalent to 40 T-75 flasks, but can support up to 2x10⁹ cells



Monoclonal Antibody Production

In the HFBR, hybridomas can yield up to 100 mg of antibody every two days, for months of culture



Recombinant Protein Production

Typical protein harvests from FiberCell Systems' cartridges contain 100 µg/mL/day or higher. Daily harvests are 2 mg to 10 mg a day.



Exosome Production

Gram quantities of exosomes can be produced in HFBRs



Cell Co-culture

The PVDF fiber allows various matrix proteins to be bound to its surface



PK/PD

Mimic human bioavailability of antibiotics



Learn more at www.fibercellsystems.com






HiSS

Diagnostics

Growth Factors

PODS® slowly release their growth factor cargo through nanopores, generating steady-state bioavailability (near zero-order kinetics)



100% protein



Slow-releasing



Versatile

PeptiGels

synthetic peptide hydrogels (PeptiGels®) and sustained release growth factors (PODS®) - to provide a reproducible and highly controlled environment for 3D cell culture



Animal-free option



Highly-cited



Reproducible

Matrix Proteins

rigorously controlled for quality and functionally tested for optimal performance in cell culture

Cell Counting

Orangu™ a non-cytotoxic, sensitive, colorimetric assay for the determination of viable cell numbers in cell proliferation and cytotoxicity assays



Non-cytotoxic



Sensitive





Easy-to-use



produced by



provided by



 www.hiss-dx.de
 hiss@hiss-dx.de


 (+49) 0761-389 49-0
 (+49) 0761-389 49-20




Cellwatcher M

Multiply your assay output by six.


Easily track cell growth, viability, motility, phenotype and drug response, without leaving your desk. Get continuous data, processed objectively and automatically, without any additional manual work steps. Benefit from higher quality of assays monitored inside the incubator and find your publication-ready results online.




Acquire multiparametric data & six conditions at a time.



Cluster growth of stem cells



Spatially resolved cell motility of a scratch



Cell motility heatmap of a scratch

Objective AI analysis, done in real-time, ensures reproducible results without the expertise, effort, and errors of manual image analysis. Simply monitor your cells' behavior across multiple conditions simultaneously and get results instantly. Top-notch visualizations and multiparametric data down to the single cell level - Cellwatcher M and PHIOme level up your analysis of e.g. wound healing, drug screens, cluster formation, and more. Access all of your data through your browser via the PHIOme cloud: anywhere, anytime.

Live cell monitoring inside the incubator. Easy.



Our Galactic Four to Monitor Your Cell Culture Processes

Countstar® Altair



Countstar® Rigel



Countstar® Mira HT



Countstar® Mira Plus



***Cell Titer – Viability – Productivity – Phenotyping
Image Based – Precise – Cost-Effective***




Cell Selection, Sorted.

Simplify and accelerate your cell line development
with Sphere Fluidics' **automated** platforms

- ✓ Completely integrated system
- ✓ Select "high producing clones" from a large transfection pool
- ✓ Cuts project lengths from months - weeks
- ✓ Dramatic reduction in reagent costs
- ✓ Provides evidence of monoclonality
- ✓ Cell friendly processing
- ✓ Traceable and certified AOF reagents and consumables
- ✓ Ease of use - low training requirements
- ✓ Small footprint






3D Bioprinting Innovations

Vital3D Technologies is developing innovative 3D bio-printing tools dedicated to the production of fully functional organs.

Company technology stands out due to its extraordinary printing efficiency thanks to resolution-on-demand capability. Our first commercially available device - Vital Light 3D - is the fastest bio-printer in the market with printable feature size down to $\sim 1 \mu$ m.





WORLD
PRECISION
INSTRUMENTS

Accelerate Your Life Science Research with Our New EVOM™ Auto High-Throughput Screening System

The Gold Standard for Automated **TEER Measurement** of Both 24 and 96 HTS Multiwell Plates



EVOM™ Auto with both 24 and 96 HTS multiwell plate capability is a high throughput screening (HTS) platform offering fast, non-invasive sample scanning by comparing electrical resistance measurements (transepithelial/transendothelial electrical resistance: TEER). TEER measurement experiments are simple to set up and less time consuming than more complex molecular studies. TEER measurement can be used as the primary scanning method to identify any physiological changes that can then be further evaluated by other methods. EVOM™ Auto can capture TEER measurements in 24 and 96 HTS multiwell plates from Corning, Millipore, or MatTek. The sample preparation time in these HTS plates is efficient, allowing for fast, multi-channel pipetting options.

APPLICATIONS



Epithelial & Endothelial
Barrier Studies




Confluence



Lung Viral Infection



Toxicity



Antibody-Antigen
Binding



Blood Brain
Barrier (BBB)



Cancer Tissue
Studies



Intestine, Kidney
and Liver Tissues

Additionally, the EVOM™ Auto electrode disinfection capability during measurement minimizes sample cross-contamination. Its wireless device control offers the convenience of operating the instrument from a distance, and the small footprint of EVOM™ Auto enables you to use it inside a sterile cell culture hood or an incubator. The EVOM™ Auto provides a fast and efficient platform for early drug discovery, by narrowing down drug targets and drug concentrations through automated, non-invasive sample scanning.

EVOM™ Auto Expedites Life Science Research by Accelerating:

- Creation of New and Effective 2-D and 3-D *In Vitro* Disease Models to Understand Disease Initiation and Progression
- Identification of Physiological Changes in Disease Models
- Quality Control of 2-D and 3-D *In Vitro* Tissues for Novel Drug Identification
- Target Screening, Identification and Verification
- Formulation Optimization & Compare Effectiveness
- Evaluate Cytotoxicity
- Novel Assay Development

WORLD PRECISION INSTRUMENTS



Determining The Feasibility of Transporting Retinal Organoids and Scaffolds Using the Cellbox.

Cellbox Solutions

The Gamm Lab utilizes stem cell technologies to investigate the cellular and molecular events during retinal differentiation and generate cells to develop advanced disease models. By understanding the behavior of these cell types, they hope to optimize strategies to delay or reverse the effects of blinding disorders such as retinitis pigmentosa and age-related macular degeneration.

Gamm Lab, University of Wisconsin-Madison.

“ Truth-in-advertising for battery life, and cell viability claims are sufficient for cross-campus transport. The Cellbox can be used to ship organoids without significant changes to the cell viability or performance.

THE PROJECT

Transporting organoids and biomaterial scaffolds can be very stressful to delicate retinal neurons and typically requires the use of special shipment media. Cryopreservation or transport at ambient temperature is often not possible due to the fragile nature of the cells and their extracellular matrix. For this reason, the Gamm Lab looked for alternatives to shipping delicate cell products and found the Cellbox.


The purpose of this trial was to evaluate the Cellbox as a viable means of transporting the fragile retinal organoids and biomaterial scaffolds. This was done in two experiments - the initial testing of the empty device; checking the cell viability and usage feasibility pre- and post-transport.

Using a controlled heating device that also

provides CO₂ allows for a safe transport of these cells under standard incubation conditions. Cellbox offers a solution to transport samples in a controlled environment with temperature and CO₂ regulated during product maturation and maintenance phases.

The Cellbox device was first tested to confirm the advertised running times. Once this was validated, organoids were packaged according to Cellbox recommendations and transported across campus to another building for imaging.

Total organoid TdTomato fluorescence (an indicator of photoreceptor survival) and cell viability tests were performed before and after the transport, and the comparison yielded no significant differences in any of the samples after Cellbox transport.



Agrostemmoside E: a potent tool for transfection assays

Transfection is the process to bring nucleic acids into eukaryotic cells. However, on the way to the cell, protection mechanisms kick in, which degrade the foreign genetic material. Hence, chemical modifications of the nucleotides and protective nanoparticles were developed. In this way, it is understood that the nanoparticle is endocytosed once it reaches the cellular membrane. Thereby, the genetic cargo reaches intracellular compartments such as early endosomes, which are acidified to become lysosomes. This process leads to further degradation and it is established that *only one percent* of the genetic material is able to escape the endosome into the cytosol, making this step **the major current obstacle** for efficient delivery of the genetic cargo.

Plant secondary metabolites, such as triterpene glycosides (saponins) have an amphiphilic structure, which allow them to interact with cellular membranes and trigger endosomal escape. In our group we could establish the continuous isolation of Agrostemmoside E (AG1856) in high purity. This saponin offers a beneficial therapeutic window and reliable, cell line dependent results. Application of this molecule to cellular assays can reduce genetic material needed for any transfection assay, since more genetic material escapes the degradation process.

Spheroid culture of murine primary chondrocytes and ATDC5 chondrocyte cell line for musculoskeletal research

Authors: Melanie Timmen, Simone Niehues, Amelie Franke, Elnaz Enderami, Richard Stange

Institute of Musculoskeletal Medicine, Department of Regenerative Musculoskeletal Medicine, University of Muenster, Muenster, Germany

Background: Cartilage tissue consists of chondrocyte cells and surrounding extracellular matrix. Chondrocytes pass through different stages from progenitor cell (origin from mesenchymal stem cells) to proliferating cells that secrete proteins forming the extracellular matrix to hypertrophic chondrocytes that form bloated cell bodies. The extracellular matrix contains different type of collagens mainly, but also a wide variety of proteoglycans, growth factors and cytokines dependent on the impact of the tissue. Cartilage appears mostly avascular and can be calcified e.g. within the growth plate or during bone healing. Growth of chondrocytes *in vitro* seems to be difficult, because of different behavior of cells in plastic associated 2D cultures or as 3D, which is often performed in micromasses or hanging drop cultures. We focused on investigation of chondrocyte growth from different sources in small spheroids and characterize the behavior of the cells. Our aim is to reconstitute cellular interaction of chondrocytes with other cell types like endothelial cells and osteoblasts.

Methods: Isolation of primary chondrocytes: The cartilaginous knee joint (hind limb) of 3-5days old mice were sequentially digested using 3mg/ml Collagenase IV. Cells were seeded in non-adherent U-bottom wells and cultured for up to 32 days in basal medium (DMEM low glucose, 10% FBS, 0,2mM L-Ascorbat-2-Phosphate, 100 U/ml Penicillin/Streptomycin (P/S)) or differentiation medium (DM) (basal medium with 100nM dexamethasone, 1mM pyruvate, 40µg/ml proline, 10ng/ml TGFbeta1, ITS-Premix. ATDC5 cells were cultured in DMEM F12, 5% FBS, 1mM pyruvate, 100U/ml P/S, and ITS-Premix. Spheroid of both cell types were seeded with 2.500 and up to 30.000 cells per well, medium was changed every 48h and cells were cultured at 37°C under normoxia (5% CO₂). For analysis, spheroids were fixed using 4% paraformaldehyde and embedded in paraffin. Slices (5µm) were stained for proteoglycans using Alcian blue staining. Immunohistocytology with antibodies against collagens as well as quantitative real time PCR to evaluate mRNA expression of marker genes for chondrocyte differentiation was performed.

Results: Primary chondrocytes build up nice spheroids with every condition used although some loose cells did not join the spheroid. Neither the two different media nor the different cell numbers/spheroid negatively affected the formation of spheres. We observed no necrotic areas within the spheroid tissue especially within the bigger spheroid or after long cultivation time (32 days). Alcian blue staining revealed the formation of extracellular matrix from day 7 on. From day 14 on cellular morphology pointed to the appearance of hypertrophic chondrocytes that was stable until day 32. However, at the outer edge of the spheroid alcian blue negative areas could be found. Collagen Type 2 positive staining was found at day 7 at the outer edge of the spheroid getting thicker with time. Using chondrocyte differentiation medium expression of marker genes like Col2a1 (collagen type II) and colX (collagen type X, hypertrophic chondrocytes) as well as Sox 9 were stimulated. ATDC5 spheroids formed faster and no loose cells were observed. Spheroid size increased during cultivation pointing to further proliferation. Alcian blue staining also revealed the formation of matrix starting at day 14, so the differentiation of chondrocytes seems to be somewhat slower compared to primary cells. Even after 21 days of culture or in bigger spheroids, no necrotic regions have been noticed. At day 21, cellular morphology indicated the onset of hypertrophic chondrocyte development. However, due of the high cell density (30,000 ATDC5 cells), alcian blue negative areas could be detected.

Conclusion: Both, primary chondrocytes and ATDC5 cell line, form nice cartilaginous spheroids with appropriate extracellular matrix, differentiation of cells from pre- to hypertrophic cell morphology and expression of marker genes. Both reveal to be a good basis for development of co-cultures with other cell types like endothelial cells and osteoprogenitors.



Next level organoid culturing

with the **RPMotion** from Orgonex

Meet the best organoid culturing device on the market

A bold claim that we're happy to back up. We have collected scientists' needs and wishes through experiments and interviews. The ultimate version of the stirring rod emerged from extensive testing by a team of engineers from Utrecht University and the Eindhoven University of Technology, the Netherlands. Orgonex' founders and their scientists, specialists in the field of organoids, experimented extensively to create the best protocols for different organoid cultures.

RPMotion

Is a semi-automated bioreactor system consisting of four 50 mL vessels and stirring rods that have been optimized for organoid culture



Strong reduction in costs

Reduce up to 75% in labor and 60% of medium expenditure

High organoid yields compared to static culturing

Proven 5-fold and up increase cellular yield

RPMotion

has a modular setup and fits into existing laboratory infrastructure

Get rid of the tedious static culturing process by using **RPMotion**

Contact us

Email: info@orgonex.com

Orgonex BV

Padualaan 8
3584 CH Utrecht
The Netherlands

Registered at the Dutch Chamber
of Commerce under no. 88954870

Get in touch

Book your demo

Are you interested in the RPMotion? Scan the QR code and fill out your details. We'll contact you to schedule a free video call. In about 30 minutes we'll talk you through the features and answer all of the questions you might have.

Scan this QR code to
book a free demo call



For over two decades, Oxford Optronix has been dedicated to ensuring reproducibility in cellular research. Here we discuss two widely utilized systems that contribute to this mission in both cell manufacturing and organoid research. Firstly, the GelCount, an automated imaging and analysis platform for colony, spheroid, and organoid counting/sizing is presented. This automation not only saves time and eradicates human bias, but also greatly improves the reproducibility of all counts across a study. Secondly, the HypoxyLab incubator and workstation is introduced with its unique approach for setting a precise oxygen concentration environment independent of altitude or atmospheric pressure changes - a critical factor affecting reproducibility of every hypoxia/physoxia based study.

Listen to our talk and discover a better way to find, analyze and isolate your most **valuable cells**, with ease and speed:

Novel automated picodroplet-based technology to streamline cell line development and cell engineering workflows with Tarun Kumar Mahto, Applications Specialist

Generation of a highly expressing, regulatory acceptable clonal cell line is the first key step in the development of a biopharmaceutical. To provide the key characteristics (Productivity, Stability, Quality, Monoclonality, and Scalability) from a cell line is challenging especially when using traditional techniques. Significant bottlenecks develop when screening for these rare cells and lengthen the cell line development workflows. Similarly, cell engineering strategies for biotherapeutics discovery frequently add costs and time. Our Cyto-Mine® platform, powered by picodroplet-based microfluidic technology, streamlines the cell line development workflow. It offers a powerful solution to the challenges of screening large numbers of cells allowing isolation of the highest value clones while ensuring monoclonality and enables a viable high-throughput cell engineering workflow.



Development of an organoid-based cell transformation assay for intestinal carcinogenicity screening of chemicals

Ziwei Wang[#], Shen Chen, Fei Chen, and Wen Chen*

Department of Toxicology, School of Public Health, Sun Yat-sen University, Guangzhou 510080, China.

Abstract The high incidence of colorectal cancer (CRC) is closely associated with the exposure to environmental carcinogens. In order to assess intestinal carcinogenic potential of chemicals, we established an adult stem cell-derived intestinal organoid model from mouse intestine (mASC-IOs) and develop a cell transformation assay (CTA). Typical chemicals were selected to assess the intestinal carcinogenesis using mASC-IOs-based CTA. The experimental design was optimized including dose range, frequency and duration of administration, indicators and period of experiment and so on. Sixteen environmental chemicals (Cd, iPb, Cr-VI, iAs-III, Zn, Cu, PFOS, BPA, PCB-126, MEHP, AOM, DMH, BaP, MNNG, Aspirin, Metformin) were selected to test the sensitivity, specificity, and accuracy of this assay. mASC-IOs maintained the biological characteristics of their origins in terms of self-organization, proliferation, differentiation, organoid morphology, genetic stability following limited splitting. To conduct mASC-IO-based CTA, mASC-IOs were treated with chemicals for 72 h after seeding, and the treatment was repeated 4 times. The carcinogenic latency of chemicals was examined for at least 10 weeks, determined by single-cell colony-formation rate in colony formation assay and anchorage-independent growth assay, and confirmed by tumorigenicity assay in nude mice. The results of carcinogenicity from 16 chemicals were highly corresponding to the epidemiological results (10/13), *in vivo* mouse studies (12/16), *in vitro* studies (11/14) and CTA results conducted following OECD guideline (6/11). Increasing proliferation of mASC-IOs was an early key event in chemical-induced oncogenic transformation. Frequency of colony formation in single-cell colony formation assay and anchorage-independent growth assay was significantly correlated with tumor formation rate in nude mice, which could be used to analyze the dose-dependent carcinogenic potential. We also found that the degree of tumor malignant in nude mice was determined by the differentiation level of mASC-IOs in the group of chemicals with carcinogenicity. In addition, the expression level of the most commonly mutated genes in human CRC, including APC, TP53, KRAS, and SMAD4 were highly correlated with the activity of chemical-induced oncogenic transformation. Taken together, we have successfully developed a mASC-IO-based CTA for intestinal carcinogenicity screening of chemicals, which plays an important role in promotion of the development and application of organ-specific screening systems.

This work was supported by NSFC 82204080 and NKPs 2017YFC1600200.

Keywords: Colorectal cancer, Intestinal organoids, Cell transformation, Environmental chemicals, Carcinogenicity screening

PeptiGels™: A technological platform for the design of fully defined synthetic 3D hydrogel scaffolds for cell engineering applications.

Prof Alberto Saiani, PhD, FRSC

Division of Pharmacy and Optometry & Manchester Institute of Biotechnology, School of Health Sciences, Faculty of Biology, Medicine and Health, The University of Manchester, UK

E-mail: a.saiani@manchester.ac.uk

Web: www.polymersandpeptides.co.uk

The use of non-covalent self-assembly has become a prominent strategy for the construction of increasingly functional biomaterials for a range of applications. A variety of molecular building blocks can be used for this purpose, one such block that has attracted considerable attention in the last 20 years is *de-novo* designed peptides. Our group work focusses on the development of a technological platform for the design of novel biofunctional 3D hydrogels scaffolds exploiting the self-assembly of so-called β -sheet forming peptides. These scaffolds can be easily functionalised using specific biological. Through the fundamental understanding of the self-assembly and gelation processes of these peptides across length scales we have been able to design hydrogels with tailored properties.








Figure: (Top) Schematic representation of the self-assembly process of β -sheet forming peptides. (Bottom) Characterisation across length scales for molecular assembly to network cross-linking. (Right) Examples of injectable, sprayable and functional hydrogel scaffolds

These 3D hydrogel scaffolds have been used in a range of contexts including cell culture, drug delivery tissue engineering, bioprinting and organoid engineering:



al. Adv. Healthcare Mat., 1900410 (2019)

vski et al. ACS Nano, 16, 4322–4337 (2022)

Videos and **publications** are available from our group [website](http://www.polymersandpeptides.co.uk).

Fraunhofer ISC – TLZ TERM

AI-powered Analysis in Robot-Assisted Three-dimensional Tissue Models Production

Dalia Mahdy, PhD Researcher

International Forum on Cell Manufacturing & Engineering
7-8.11.2023. Berlin, Germany

Creation of skin cell spheres and improved methods for measuring cell viability in the 3D model

Iryna Levkovych¹, Adrianna Maria Piasek, MSc.Eng.¹, Michał Stepulak, Ph.D.², Anna Sobiepanek, Ph.D.^{1,*}

¹Laboratory of Biomolecular Interaction Studies, Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warsaw

²BASF Poland Sp. z o.o., al. Jerozolimskie 142B, 02-305 Warsaw

*Corresponding author: anna.sobiepanek@pw.edu.pl

The most common *in vitro* model of skin for testing pharmaceuticals and cosmetic products is the 2D cell monolayer. However, it has a significant drawback as it oversimplifies the interactions within the model (1). In contrast, 3D *in vitro* models, including spheres and equivalents, provide a more accurate representation of cell-cell interactions. Several methods for producing spheres, such as the hanging drop method (HD) and limiting cell adhesion (LCA), are widely known (2-3). Unfortunately, they usually require additional equipment or do not provide sufficient repeatability (4). Moreover, many existing protocols for cell viability assays are tailored to 2D models, hence the optimization step before conducting experiments on the 3D *in vitro* models is needed. Therefore, our study aims to optimize methods of sphere creation and to adjust the existing viability assays for their usage in these advanced models.

This study was conducted on the primary human skin cells and commercial cell lines. The HD and the LCA methods were compared in the context of sphere formation. To determine which surfactant would better limit the cell adhesion Pluronic F127 (Poloxamer 407), Pluronic F68 (Poloxamer 188) and Kolliphor EL (PEG35 castor oil) were applied as coating agents. Cell viability was analyzed by the dehydrogenase (MTT) and esterase (FDA) activity assays.

The LCA method provided more repeatable results of the sphere formation than the HD method. Above all, spheres differed in morphology depending on the used cell type. Different melanoma cell lines showed varied capacity to form compact cell aggregates, which could be correlated to their adhesion properties visible in the 2D culture. A low cytotoxic influence of the two coating agents Pluronic F127 and F68 on skin cells was confirmed, whereas the application of Kolliphor EL led to cell membrane damage. The surfactant's influence on the cell adhesion depended on its concentration. The influence of the extracellular matrix proteins on the process of cell arrangement into aggregates is minor. Although the FDA assay can be directly applied to the sphere model, the MTT protocol needs modifications to improve the dissolvent of formazan from cell spheres.

Our findings indicate that the basic 3D cell culture protocols can be further optimized to facilitate the preparation of these models and their usage.

This study was supported by the Dean's grant (2023) from the subsidy of the Faculty of Chemistry of the Warsaw University of Technology.

Keywords: 3D *in vitro* models, skin cells, spheres, limiting cell adhesion, surfactants.

1. Piasek et al. 2023, Journal of Visualized Experiments 200, e65773.
2. Lv et al. 2017, Oncology Letters 14 (6), 6999–7010.
3. Costa et al. 2018, Biotechnology Journal 13:1700417.
4. Amaral et al. 2017, Frontiers in Physiology 8, 605.

Empowering Breakthroughs for Life.

See You in 2024

5-6 November 2024

Berlin, Germany

